

VIRO-STAT

Disinfectant Cleaner

Form: Liquid

Color: Colorless, Clear

Odor: Mild

pH: 12.40

Solubility (in water): Soluble
(in mineral spirits): No data available

VOC Content (% by weight): <1.0%

Flash Point (Closed Cup): >200 °F

Specific Gravity: 1 g/cm³

Density: 8.42 lbs /gal

<u>Ingredients</u>	<u>C.A.S. #</u>
Water	7732-18-5
Alcohol ethoxylate.....	68131-39-5
Quaternary ammonium compounds, benzyl- C12-16-alkyldimethyl, chlorides.....	68424-85-1
Tetrasodium EDTA	64-02-8
Decyldimethyloctylammonium chloride	32426-11-2

Ideal for...

- Restrooms
- Locker Rooms/Shower Stalls
- Playground Equipment
- Institutional Kitchens
- Hospitals/Patient Care Facilities
- Operating Rooms
- Day Care Centers
- Veterinary Clinics/Kennels
- Farms/Hatcheries/Aviaries
- Sewer Backup/River Flooding

Effective against...

- E. coli
- Staphylococcus aureus
- MRSA & VISA
- HBV & HCV
- HIV-1

and many more...

DIRECTIONS:

Preparation of Disinfectant Use-Solution: Add (mix) 2 oz. per gallon of water (1:64). For heavy duty use, add (mix) 8 oz. per gallon of water (1:16). Add 8 oz. per gallon to kill Adenovirus Type 5 and Type 7.

Disinfection/Virucidal/Fungicidal/Mold and Mildew Control Directions: Apply use-solution to hard, nonporous surfaces, thoroughly wetting the surfaces, with a cloth, mop, sponge, sprayer or by immersion. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface; rub with brush, sponge or cloth.

Veterinary Practice/Animal Care Disinfection Directions: For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc.. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floor, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 2 oz. of VIRO-STAT per gallon of water (or equivalent dilution) for a period of 10 minutes. Wipe or allow to air dry. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and manure. Rinse all surfaces that come into contact with food, including equipment used for feeding or watering, with potable water before reuse. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

CONSULT PRODUCT LABEL FOR DETAILED USE INSTRUCTIONS

HMIS®				NFPA®	
		Severe	4	Extreme	
Health	3	Serious	3	High	Health..... 3
Flammability.....	1	Moderate	2	Moderate	Flammability 1
Reactivity.....	0	Slight	1	Slight	Reactivity 0
Personal Protection	X	Minimal	0	Insignificant	Special PrecautionsNone